



a-c, e (independently selected) = 0 or 1;
d = 0;
R = modifying group, sialyl or oligosialyl

FIG. 29A



FIG. 29B



FIG. 29C



FIG. 29D



FIG. 29E



FIG. 29F



FIG. 29G





a-d, i, n-u (independently selected) = 0 or 1.
aa, bb, cc, dd, ee (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 20.
v-z = 0; R = modifying group, mannose, oligo-mannose.
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 30A



FIG. 30B



FIG. 30C


Yeast expressed interferon alpha-14C.
a-q, cc, dd, ee, v-z = 0;
r-y (independently selected) = 0 to 1;
aa, bb = 1;
R (branched or linear) = Man, oligomannose or polysaccharide.
1. Endo-H
2. Galactosyltransferase, UDP-Gal
3.. CMP-SA-PEG, ST3Gal3
a-z, bb = 0; aa = 1; R' = -Gal-Sia-PEG.

FIG. 30D





a-d, i, r-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 or 1.
n, v-y = 0; z = 0 or 1.
R = polymer; R' = sugar, glycoconjugate.

FIG. 30E



FIG. 30F



FIG. 30G



FIG. 30H



FIG. 30I



FIG. 30J



FIG. 30K



FIG. 30L



FIG. 30M


Insect cell or fungi expressed interferon alpha-14C.
a-d, f, h, j-n, s, u, v-y = 0;
e, g, i, r, t (independently selected) = 0 or 1; z = 1.
1. endoglycanase
2. Galactosyltransferase,
UDP-Gal-linker-SA-CMP
3. ST3Gal3, transferrin
i (independently selected) = 0 or 1;
a-h, j-m, r-z = 0;
n = 1; R' = -Gal-linker-transferrin.

FIG. 30N





a-c, e, f (independently selected) = 0 or 1;
d, g = 0; R = polymer, glycoconjugate.

FIG. 30O



FIG. 30P



FIG. 30Q



FIG. 30R



FIG. 30S



FIG. 30T



FIG. 30U



FIG. 30V



FIG. 30W









a-c, e (independently selected) = 0 or 1;
d = 0; R = polymer, glycoconjugate.

FIG. 30X



FIG. 30Y



FIG. 30Z


E. coli expressed interferon alpha-mucin fusion protein.
a-e = 0.
1. GalNAc Transferase, UDP-GalNAc
2. CMP-SA-PEG, sialyltransferase
c, d, e (independently selected) = 0 or 1;
a, b = 0; R = PEG.

FIG. 30AA











a-c, e (independently selected) = 0 or 1;
d = 0; R = polymer, linker.

FIG. 30BB



FIG. 30CC



FIG. 30DD


E. coli expressed Interferon alpha (no fusion).
a-e, n = 0.
1. NHS-CO-linker-SA-CMP
2. ST3Gal3, transferrin
a-e = 0; n = 1; R' = linker-transferrin.

FIG. 30EE





a-d, i, r-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 or 1.
n, v-y = 0; z = 0 or 1; R = polymer

FIG. 31A



FIG. 31B



FIG. 31C



FIG. 31D



FIG. 31E



FIG. 31F



FIG. 31G



FIG. 31H



FIG. 31I



FIG. 31J



FIG. 31K

NSO expressed Ifn-beta.
a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n, v-y = 0;
Sia (independently selected) = Sia or Gal.

1. Sialidase and α-galactosidase
2. α-Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3

a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; R = PEG
n = 0; v-y (independently selected) = 1,
when j-m (independently selected) is 1;

FIG. 31L

CHO, BHK, 293 cells, Vero expressed Ifn-beta.
a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n, v-y = 0.

1. Sialidase
2. CMP-SA-PEG (16 mol eq), ST3Gal3
3. CMP-SA, ST3Gal3

a-d, i-m, r-u, z (independently selected) = 0 or 1;
e-h = 1; n = 0;
v-y (independently selected) = 0 or 1; R = PEG.

FIG. 31M



FIG. 31N



FIG. 31O





a-d, i, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = modifying group;
R' = H, glycosyl group, modifying group, glycoconjugate.

FIG. 31P



FIG. 31Q



FIG. 31R


CHO, BHK, 293 cells, Vero expressed Ifn-beta.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-SA-CMP, ST3Gal3
2. ST3Gal3, desialylated transferrin.
3. CMP-SA, ST3Gal3
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-transferrin.

FIG. 31S





B ◄ {Glc-(Xyl)n }o

C ◄ {-Fuc }p

a-d, i, q-u (independently selected) = 0 or 1.
o, p (independently selected) = 0 or 1.
e-h, n (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 20.
v-y = 0;
R = modifying group, mannose, oligo-mannose, Sia-Lewis X, Sia-Lewis A..

FIG. 32A



FIG. 32B



FIG. 32C


NSO expressed Factor VII or VIIa
a--u (independently selected) = 0 or 1;
v-y = 0; n = 0-4;
Sia (independently selected) = Sia or Gal.
1. Sialidase and α-galactosidase
2. Galactosyltransferase, UDP-Gal
3. CMP-SA-PEG, ST3Gal3
a-m, o-u (independently selected) = 0 or 1;
n = 0-4; v-y (independently selected) = 1,
when j-m (independently selected) is 1;
Sia = Sia; R = PEG.

FIG. 32D







a-d, i, n-u (independently selected) = 0 or 1.
bb, cc, dd, ee, ff, gg (independently selected) = 0 or 1.
e-h, aa (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 20.
v-z = 0; R = modifying group, mannose, oligo-mannose.

FIG. 33A



FIG. 33B



FIG. 33C



FIG. 33D



FIG. 33E



FIG. 33F



FIG. 33G





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose.

FIG. 34A



FIG. 34B



FIG. 34C



FIG. 34D



FIG. 34E



FIG. 34F



FIG. 34G



FIG. 34H



FIG. 34I


CHO, BHK, 293 cells, Vero expressed FSH.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-SA-CMP, ST3Gal3
2. ST3Gal1, desialylated chorionic gonadrophin (CG) produced in CHO.
3. CMP-SA, ST3Gal3, ST3Gal1
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-CG.

FIG. 34J







a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0;
R = polymer.

FIG. 35A



FIG. 35B



FIG. 35C



FIG. 35D



FIG. 35E



FIG. 35F



FIG. 35G



FIG. 35H



FIG. 35I



FIG. 35J



FIG. 35K



FIG. 35L



FIG. 35M



FIG. 35N



FIG. 35O



FIG. 35P



FIG. 35Q



FIG. 35R



FIG. 35S





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0; aa, bb = 1; cc = 0;
R = polymer; R'' and R' = sugar-polymer or Fuc.

FIG. 35T



FIG. 35U



FIG. 35V


Insect cell expressed EPO
a-d, f, h, j-m, n-q, s, u, v-z = 0;
e, g, i, r, t (independently selected) = 0 or 1;
aa = 1; R" = Fuc.
1. mannosidases
2. galactosyltransferase, UDP-Gal-PEG
cc, e, i, r, v (independently selected) = 0 or 1;
bb, a-h, j-q, s-u, w-z = 0; aa = 1; R' = Gal-PEG.

FIG. 35W







a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
v-z = 0;
R = polymer.

FIG. 35X



FIG. 35Y



FIG. 35Z


CHO, BHK, 293 cells, Vero expressed NESP
a-g, n, q = 1; h = 1 to 3;
j-m, i, o, p (independently selected) = 0 or 1;
r-u (independently selected) = 0 to1; v-z = 0
1 CMP-SA, poly-α2,8-ST
a-g, n, q = 1; h = 1 to 3;
i, j-m, o, p, r-u, (independently selected) = 0 or 1;
v-z (independently selected) = 0-40; R = Sia.

FIG. 35AA







a-d, i, n-u, aa (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer, glycoconjugate.

FIG. 36A



FIG. 36B



FIG. 36C



FIG. 36D



FIG. 36E



FIG. 36F



FIG. 36G







a-d, i, n-u, aa, bb, cc (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = modifying group, mannose, oligo-mannose.
R'= H, glycosyl residue, modifying group. glycoconjugate.

FIG. 36H



FIG. 36I



FIG. 36J


CHO, BHK, 293 cells, Vero expressed GM-CSF.
a--m, o-u, aa, bb (independently selected) = 0 or 1;
n, v-z, cc = 0.
1. sialidase
2. CMP-SA, ST3Gal3
2. CMP-SA-linker-SA-CMP, ST3Gal1
3. ST3Gal3, transferrin
a--m, p-u, z, aa (independently selected) = 0 or 1;
o, v-y, cc = 0; bb, n = 1; R = transferrin.

FIG. 36K





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer.

FIG. 37A



FIG. 37B



FIG. 37C



FIG. 37D



FIG. 37E



FIG. 37F



FIG. 37G





a-d, i, n, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose;
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 37H



FIG. 37I



FIG. 37J



FIG. 37K



FIG. 37L



FIG. 37M



FIG. 37N





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer.

FIG. 38A



FIG. 38B



FIG. 38C



FIG. 38D



FIG. 38E


CHO, BHK, 293 cells, Vero expressed α_1-antitrypsin.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA, α2,8-ST
a-d, i, q-u (independently selected) = 0 or 1; e-h = 1;
j-m (independently selected) = 0-20;
v-y (independently selected) = 0.

FIG. 38F





a-d, i, n, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose;
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 38G



FIG. 38H



FIG. 38I


CHO, BHK, 293 cells, Vero expressed α_1-antitrypsin.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-Gal-UDP, ST3Gal3
2. Galactosyltransferase, transferrin treated with endoglycanase
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0;
v-y (independently selected) = 0 or 1;
R = linker-transferrin.

FIG. 38J





a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
R = polymer;
R', R" (independently selected) = sugar, glycoconjugate.

FIG. 38K



FIG. 38L



FIG. 38M


CHO, BHK, 293 cells, Vero, transgenic animal expressed α_1 antitrypsin.
a-h, i-o, r-u (independently selected) = 0 or 1;
p, q, v-y = 0.
1. CMP-SA-PEG, ST3Gal3
a-h, i-o, r-u (independently selected) = 0 or 1;
p, q = 0; v-y (independently selected) = 0 or 1;
R = PEG.

FIG. 38N





a-d, i, q-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0; R = polymer.

FIG. 39A



FIG. 39B



FIG. 39C



FIG. 39D



FIG. 39E


CHO, BHK, 293 cells, Vero expressed Cerezyme.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA, α2,8-ST
a-d, i, q-u (independently selected) = 0 or 1;
e-h = 1; j-m (independently selected) = 0-20;
v-y (independently selected) = 0.

FIG. 39F





a-d, i, n, p-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 6.
j-m (independently selected) = 0 to 100.
v-y = 0;
R = modifying group, mannose, oligo-mannose;
R' = H, glycosyl residue, modifying group, glycoconjugate.

FIG. 39G



FIG. 39H



FIG. 39I


CHO, BHK, 293 cells, Vero expressed Cerezyme.
a-d, i-m, q-u (independently selected) = 0 or 1;
e-h = 1; v-y = 0.
1. CMP-SA-linker-SA-CMP, ST3Gal3
2. ST3Gal3, desialylated transferrin.
3. CMP-SA, ST3Gal3
a-m, q-u (independently selected) = 0 or 1;
p = 1; n = 0; v-y (independently selected) = 0 or 1;
R = linker-transferrin.

FIG. 39J







a-d, i, n-u (independently selected) = 0 or 1.
e-h (independently selected) = 0 to 4.
j-m (independently selected) = 0 to 20.
R = polymer; R' = sugar, glycoconjugate.

FIG. 40A



FIG. 40B



FIG. 40C



FIG. 40D



FIG. 40E



FIG. 40F



FIG. 40G